

The expectation value of the metric operator with respect to Gaussian weave state in loop quantum gravity

Tomoya Tsushima*

Institute for Theoretical Physics, Kanazawa University, Kanazawa 920-1192, JAPAN

(Dated: February 4, 2008)

Abstract

Loop Quantum Gravity is the major candidate of quantum gravity. It is interesting to consider its continuum limit, which corresponds to the classical limit. We consider the Gaussian weave state, which describes a semi-classical picture. We calculate the expectation value of metric operator with respect to this state.

arXiv:gr-qc/0212117v3 7 Nov 2003

*Electronic address: tomoya@hep.s.kanazawa-u.ac.jp

I. INTRODUCTION

In the twenty years, the canonical quantum gravity has made great progress [1]. We call this new stage of theory the loop quantum gravity (LQG). The quantum state is characterized by closed paths on three-dimensional space, and is called spin network state. Its norm is positive definite. It has been proved non-perturbatively that the spin network state is an eigen state with respect to three-dimensional geometrical observables such as area and volume, its eigen value is discrete [2, 3, 4, 5, 6]. The black hole entropy is calculated by counting the paths of quantum states crossing the surface of the (classical) event horizon and is proportional to the area eigenvalue of the surface [7]. The non-commutativity of the spatial observable has been also argued in this framework [8]. Note that the Hamiltonian constraint operator, which is strongly related to dynamics of the system, is not been solved so far. Although the matrix element of it has been calculated with respect to the spin network state [9, 10].

Recently, the ultra high energy cosmic ray (UHECR) events, whose energy is larger than 10^{11} GeV, have been detected by AGASA experiment [11, 12]. Assume that the UHECR is proton, it can not travel more than 10^2 Mpc because it interacts with the cosmic microwave background radiation. Since there is no nearby sources in our galaxy, the charged particle cosmic ray spectrum must obey a high energy energy cutoff, which is called Greisen-Zatsepin-Kuzmin (GZK) cutoff [13, 14]. However, it conflicts to the data from AGASA. Then, the AGASA data is a great challenge.

This problem can be solved by taking account of the possible quantum gravity effects deforming the dispersion relation $E^2 = \vec{p}^2 + m^2$ due to the Lorentz invariance [15, 16]. It is nothing that violation of the relativity first induced by [17] with spontaneous breaking down mechanism. In fact, the semi-classical approximation of quantum gravity generally derives

$$E^2 = \vec{p}^2 + m^2 + \sum \xi_n |\vec{p}|^n / M_{\text{Pl}}^{n-2} \quad (1)$$

where coefficients ξ_n 's are matter dependent parameters. In LQG, this relation is derived by [18, 19, 20, 21]. Roughly speaking, LQG is a quantum mechanical system in the space of $(SU(2))^n$. The wave functional is constructed by n holonomies, and is square integrable with direct product of n $SU(2)$ Haar measures. A holonomy is a path-ordered exponential of $SU(2)$ connection. In semi-classical approximation, the expectation value of the connection is required as a classical variable instead of the holonomy. If the holonomy is expanded in

power of the connection, or of the length of the holonomy, then spatially non-local terms are generated. Lorentz symmetry violates. The mystery of UHECR has been also approached by enlarging the standard model of elementary particles without deforming the dispersion relation [22, 23, 24, 25, 26].

More concretely, let us consider the (densitized inverse) dreibein operator ${}^n\hat{\Theta}_I^i$, which plays a role of gravitational part of Hamiltonian for matter field. It is expanded with respect to the length of holonomy as

$$\begin{aligned} {}^n\hat{\Theta}_I^i &= \frac{4}{i\hbar\kappa} \text{tr}(\tau^i h_I \hat{V}^n h_I^{-1}) \\ &\equiv \frac{2}{i\hbar\kappa} \dot{s}_I^a [\hat{A}_a^i, \hat{V}^n] - \frac{1}{i\hbar\kappa} \dot{s}_I^a \dot{s}_I^b [\partial_a \hat{A}_b^i + \frac{1}{2} \epsilon^{ijk} \hat{A}_a^j \hat{A}_b^k, \hat{V}^n] + \mathcal{O}(|\dot{s}_I^a|^3), \end{aligned} \quad (2)$$

where $\hat{V}$ is the volume operator. The expectation value of it is estimated in terms of the orders of $\langle \cdots \hat{A}_a^i \cdots \rangle \equiv \cdots \times (\ell_{\text{Pl}}/\mathcal{L})^\Upsilon / \mathcal{L} \times \cdots$, where $\mathcal{L}$ is a scale that the space looks like continuum for fields. Because $\hat{A}$ is not defined and state is not decided, parameter Υ is totally unknown and the direction and the length of $\dot{s}^a$ are also unknown. These ambiguous situation comes from the lack of information about the quantum system.

Another discussion, the coherent state [29] is adopted as a semiclassical state, and is constructed the cubic graph states [27, 28]. The state has a dimensionless parameter, which can be smaller than one, the semi-classical approximation is characterized by this parameter. The cubic graph is complicated because its vertex is hexavalent. (The “valence” is defined in the next section.) For weak gravity and calculation simplicity, the gauge group is reduced from $SU(2)$ to $U(1)^3$.

In this paper, we employ a Gaussian weave state [30], which is superposition of infinite number of states so that it can describe semiclassical space. Then we calculate the dreibein operator before series expansion. Furthermore, the metric operator, sum of squared dreibein operator, is calculated in order to determine a mean value of $\dot{s}^a$. We consider the purely gravitational part, matter contributions are not treated. In the next section, we explain a essence of LQG we need, and determine a notation. In section III, we introduce a Gaussian weave state, and calculate the expectation value of the metric operator. The techniques for spin network calculation is in appendix.

II. LOOP QUANTUM GRAVITY

A. Real Ashtekar variable

In LQG, the Einstein-Hilbert action

$$S = \frac{1}{16\pi G} \int dt d^3x N \sqrt{q} (R - (K_a^a)^2 + K_{ab}K^{ab}) \quad (3)$$

is employed in order to describe the gravitational field. $a, b, \dots$ are spatial indices and q_{ab} is a three-dimensional metric and $q = \det(q_{ab})$. R and K_{ab} are a three-dimensional Ricci scalar and an extrinsic curvature, respectively. G is the Newton constant. The lapse function N corresponds to time-time component of four-dimensional metric. Independent variables of this action are, naively, q_{ab} and its canonical momentum. Let us introduce a dreibein θ_a^i satisfies the relation $q_{ab} = \theta_a^i \theta_b^i$. The new canonical variables are a dreibein of density weight one $E_i^a = \frac{1}{2} \epsilon^{abc} \epsilon_{ijk} \theta_b^j \theta_c^k$ and a real Ashtekar variable

$$A_a^i = \Gamma_a^i + \frac{\gamma}{\sqrt{q}} E_i^b K_{ab}, \quad (4)$$

where Γ_a^i is a spin-connection that is a function of E_j^b satisfies the torsionless condition $\partial_a E_i^a + \epsilon^{ijk} \Gamma_a^j E_k^a = 0$. The Immirzi parameter γ is a real number, which cannot be determined in the theoretical point of view. The indices $i, j, \dots$ are degrees of freedom of a local internal $SO(3)$. This canonical pair forms a Poisson bracket

$$\{A_a^i(x), E_j^b(y)\}_P = \kappa \delta_a^b \delta_j^i \delta^3(x, y), \quad (5)$$

where $\kappa = 8\pi\gamma G$, which reproduce the original one that made by metric and its conjugate.

The real Ashtekar variable A_a^i behaves as an $SU(2)$ connection form. That is, we can regard A_a^i and E_i^a as a vector potential and an electric field in the $SU(2)$ gauge theory, respectively. The constraints, which included by this system, are Hamiltonian constraint

$$\mathcal{H} = \frac{\gamma}{2\kappa\sqrt{q}} \epsilon^{ijk} E_i^a E_j^b (F_{ab}^k - (\gamma^2 + 1) \epsilon^{klm} K_a^l K_b^m), \quad (6)$$

Gauss constraint $\mathcal{G}^i = \frac{1}{\kappa} (\partial_a E_i^a + \epsilon^{ijk} A_a^j E_k^a)$ and diffeomorphism constraint $\mathcal{D}_a = \frac{1}{\kappa} E_i^b F_{ab}^i - A_a^i \mathcal{G}^i$, where $F_{ab}^i = \partial_a A_b^i - \partial_b A_a^i + \epsilon^{ijk} A_a^j A_b^k$ is a curvature 2-form of A_a^i , and $K_a^i = \frac{1}{\gamma} (A_a^i - \Gamma_a^i)$. $\mathcal{H}$, $\mathcal{G}^i$ and $\mathcal{D}_a$ are caused by time reparametrization, $SU(2)$ gauge transformation and spatial diffeomorphism invariance, respectively.

B. Regularization

In (6), the weight $\sqrt{q}$ locates a denominator, because the canonical momentum E_i^a is density weight one and the integrand must be density weight one. As an example, the electromagnetic Hamiltonian is

$$H_{EM}[N] = \int d^3x \frac{1}{2} N \frac{q_{ab}}{\sqrt{q}} (\underline{E}^a \underline{E}^b + \underline{B}^a \underline{B}^b). \quad (7)$$

Since both electric and magnetic field $\underline{E}^a, \underline{B}^a$ are density weight one, the integrand also has an inverse weight. If we naively quantize this, a second order functional derivative emerges at the same point and the gravitational sector diverges. Therefore this operator is ill-defined. However, using volume variables and point splitting methods, we can solve these two types of singularities [31, 32].

Consider a box, which satisfies $\int_{\text{Box}} d^3y = \epsilon$, centered at x , and $f_\epsilon(x, y)$ is unity if y in the box and is zero otherwise. If we take a limit $\epsilon \rightarrow 0$, then $\frac{1}{\epsilon} f_\epsilon(x, y) \rightarrow \delta^3(x, y)$. The volume variable in the box is

$$\begin{aligned} V_\epsilon(x) &= \int d^3y f_\epsilon(x, y) \sqrt{q(y)} \\ &= \int d^3y f_\epsilon(x, y) \sqrt{\left| \frac{1}{6} \epsilon_{abc} \epsilon^{ijk} E_i^a E_j^b E_k^c \right| (y)} \end{aligned} \quad (8)$$

This becomes $\frac{1}{\epsilon} V_\epsilon(x) \rightarrow \sqrt{g(x)}$ as $\epsilon \rightarrow 0$. Poisson bracket of A_a^i and V^n becomes a dreibein of density weight $(n-1)$:

$$\left(\frac{\theta_a^i}{(\sqrt{q})^{1-n}} \right) (x) = \lim_{\epsilon \rightarrow 0} \epsilon^{1-n} \frac{2}{\kappa n} \{A_a^i(x), (V_\epsilon(x))^n\}_P \quad (9)$$

In particular, the density weight becomes a negative if $n < 1$. Let this relation apply to (7). This is in the case of $n = 1/2$, and there is a factor $\epsilon^{1/2} \epsilon^{1/2} = \epsilon$ in the numerator. Then we should insert a point splitting $\int d^3y \frac{1}{\epsilon} f_\epsilon(x, y)$ to eliminate the ϵ . Thus,

$$\begin{aligned} \int d^3x \frac{q_{ab}}{\sqrt{q}} X^a Y^b &= \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \int d^3x \int d^3y f_\epsilon(x, y) \left(\frac{\theta_a^i X^a}{(\sqrt{q})^{1/2}} \right) (x) \cdot \left(\frac{\theta_b^j Y^b}{(\sqrt{q})^{1/2}} \right) (y) \\ &= \frac{16}{\kappa^2} \lim_{\epsilon \rightarrow 0} \int d^3x \int d^3y f_\epsilon(x, y) \\ &\quad \times \left(\{A_a^i, \sqrt{V_\epsilon}\}_P X^a \right) (x) \left(\{A_b^j, \sqrt{V_\epsilon}\}_P Y^b \right) (y) \end{aligned} \quad (10)$$

This corresponds to the regularized electromagnetic Hamiltonian. Similarly, The gravitational Hamiltonian (constraint) can be explained by the commutator with volume variable and point splitting.

We divide a spatial integration into regions specified by ϵ . We make the region at most includes one vertex of quantum states, which defined in next subsection. By this division, if we quantize a regularized variable, that becomes well-defined operator independently ϵ . Thus, the limit will be eliminated.

C. Quantum states

The $SU(2)$ holonomy is a path ordered integral of the connection form along the smooth path e ,

$$h_e^{(p)}(A) = \mathcal{P} \exp \left(- \int_0^1 ds \dot{e}^a(s) A_a^i(e(s)) \tau_{(p)}^i \right) \quad (11)$$

on the three dimensional space. We call the path e an edge, and parametrize the orbit e^a of the edge by $s \in [0, 1]$. $\dot{e}^a(s)$ is a tangent vector of the edge. Each of beginning point $e^a(0)$ and finalpoint $e^a(1)$ is called vertex. $\tau_{(p)}^i$ is an anti-hermite generator of $su(2)$ $(p+1)$ -dimensional representation, or, equivalently, spin- $p/2$ representation. This integer p is called a color of the edge. The holonomy changes by gauge transformation from h_e to $g(e(1))h_e g^{-1}(e(0))$ for $g(e(1)), g(e(0)) \in SU(2)$.

Quantum state is characterized by closed graph γ constructed by edges. Edges meet at a vertex. They don't have to connect smoothly. A vertex which connects n pieces of edges is called n -valent vertex. In trivalent vertex, for example, if colors of edges are a, b and c , respectively and if a and b are given, c can only take $|a - b|, |a - b| + 2, \dots, a + b - 2$ and $a + b$, because of $SU(2)$ invariance of quantum states. $SU(2)$ invariant multi-valent vertex can be constructed by set of trivalent vertices. Using $(n - 3)$ pieces of 'virtual' edge with gauge invariant set of colors $\vec{i} = \{i_2, i_3 \dots, i_{n-2}\}$, we can compose the n -valent vertex by $(n - 2)$ 'virtual' trivalent vertices. The virtual edges have many degrees of freedom with respect to its colors, we regard as basis of the vertex. The way of connection between virtual edges is not unique, but if we fix the basis, the other way of connection can be described by linear combination of the basis we chose. This basis, the way of connection between edges, is called the intertwiner (or intertwining tensor.) In other words, the intertwiner is a map from tensor product of incoming edges $\bigotimes_{e_I(1) \cap v} (p(e_I) + 1)$ to tensor product of outgoing edges $\bigotimes_{e_I(0) \cap v} (p(e_I) + 1)$. FIG. 1 shows pentavalent vertex, for example.

The spin network $\{\gamma, \vec{p}, \vec{\phi}\}$ that is a set of a graph γ , colors $\vec{p} = \{p(e_1), \dots, p(e_n)\}$ of




FIG. 1: An example of pentavalent vertex. Five pieces of edges with color $P_0, \dots, P_4$ have connected at the vertex. The pentavalent vertex constructed by three virtual trivalent vertices with two virtual edges of color i_2, i_3 .

edges and intertwiner $\vec{\phi} = \{\phi_{i_1}^\tau(v_1), \dots, \phi_{i_m}^\tau(v_m)\}$ of vertices $\vec{v} = \{v_1, \dots, v_m\}$. It describes a quantum state called a spin network state. The wave function is explained by

$$\psi_{\{\gamma, \vec{p}, \vec{\phi}\}}(A) = (h_{e_1} \otimes \dots \otimes h_{e_n}) \cdot (\phi_{i_1}^\tau(v_1) \otimes \dots \otimes \phi_{i_m}^\tau(v_m)). \quad (12)$$

Now we define inner product of states of $\{\gamma_1, \vec{p}_1, \vec{\phi}_1\}$ and $\{\gamma_2, \vec{p}_2, \vec{\phi}_2\}$. There is a larger graph $\gamma = \cup_{I=1}^n e_I \supset \gamma_1 \cup \gamma_2$, then inner product of states can be explained for $\psi_{\{\gamma_1, \vec{p}, \vec{\phi}\}}$ and $\psi_{\{\gamma_2, \vec{p}, \vec{\phi}\}}$ on the graph γ as

$$\begin{aligned} \langle \{\gamma_1, \vec{p}, \vec{\phi}\} | \{\gamma_2, \vec{p}, \vec{\phi}\} \rangle &= \int_{(SU(2))^n} d\mu(h_{e_1}) \dots d\mu(h_{e_n}) \overline{f_\gamma(h_{e_1}, \dots, h_{e_n})} g_\gamma(h_{e_1}, \dots, h_{e_n}) \\ &= \delta_{\gamma_1, \gamma_2} \left(\prod_{e \in \gamma} \frac{\delta_{p_1(e), p_2(e)}}{\Delta_{p_1(e)}} \right) \cdot \left(\prod_{v \in \gamma} (\phi_1(v), \phi_2(v)) \right) \end{aligned} \quad (13)$$

with the inner product of intertwiners

$$(\phi_1(v), \phi_2(v)) = \left(\prod_{\tilde{e}} \frac{\delta_{p_1(\tilde{e}), p_2(\tilde{e})}}{\Delta_{p_1(\tilde{e})}} \right) \left(\prod_{\tilde{v}} \theta(p_1(e_{1\tilde{v}}), p_1(e_{2\tilde{v}}), p_1(e_{3\tilde{v}})) \right), \quad (14)$$

where $d\mu$ is $SU(2)$ Haar measure, which normalized $\int_{SU(2)} d\mu = 1$. $\tilde{e}, \tilde{v}$ mean a virtual edge and a virtual trivalent vertex at vertex v , respectively. $e_{1\tilde{v}}, e_{2\tilde{v}}, e_{3\tilde{v}}$ are edges (or virtual ones) that connected with the virtual vertex $\tilde{v}$. The symmetrizer Δ_a and the θ -net $\theta(a, b, c)$ are defined by (A9, A10).

For simplicity, we denote a spin network state as ψ_γ . The holonomy h_s along to the segment s , is just a product operator that operates to ψ_γ adding the edge to the graph, i.e., $h_s \psi_\gamma = \psi_{s \cup \gamma}$. The canonical conjugate operator of it is a left (right) invariant vector

$(h_e \tau^i)^A_B \partial / \partial (h_e)^A_B$ ($(\tau^i h_e)^A_B \partial / \partial (h_e)^A_B$.) It connects τ^i to the vertex of the edge in the graph. These operators can be interpreted as acting on vertices. Therefore, we set aim to the vertex, and investigate the intertwiner on it. The normalized intertwiner $\tilde{\phi}_{\vec{i}}$ of the n -valent vertex can be written graphically as

$$\tilde{\phi}_{i_2, \dots, i_{n-2}}(P_0, \dots, P_{n-1}) = N_i(P_0, \dots, P_{n-1}) \begin{array}{ccccccc} & (e_0) & (e_1) & (e_2) & & (e_{n-2}) & (e_{n-1}) \\ & P_0 & P_1 & P_2 & \cdots & P_{n-2} & P_{n-1} \\ & \downarrow & \downarrow & \downarrow & & \downarrow & \downarrow \\ & \text{---} & \bullet & \text{---} & \bullet & \text{---} & \bullet \\ & & i_2 & & & & i_{n-2} \end{array} \quad (15)$$

with the normalization factor

$$N_i(P_0, \dots, P_{n-1}) = \sqrt{\frac{\prod_{x=2}^{n-2} \Delta_{i_x}}{\prod_{x=1}^{n-2} \theta(i_x, P_x, i_{x+1})}}. \quad (16)$$

where $i_1 = P_0$, $i_{n-1} = P_{n-1}$. $e_0, \dots, e_{n-1}$ are edges connecting at the vertex, and $P_0, \dots, P_{n-1}$ are its colors respectively. The inner product (14) of $\tilde{\phi}_i$ are given graphically,

$$(\tilde{\phi}_{\vec{i}}, \tilde{\phi}_{\vec{k}}) = N_{\vec{i}} N_{\vec{k}} \left(\begin{array}{c} \text{Diagram with nodes } k_2, k_{n-2}, i_2, i_{n-2} \text{ and regions } P_0, P_1, \dots, P_{n-2}, P_{n-1} \end{array} \right) = \prod_{x=2}^{n-2} \delta_{i_x}^{k_x}. \quad (17)$$

If an operator $\hat{X}$ acts to it, we obtain a matrix element $X_i^{\vec{k}} = (\tilde{\phi}_{\vec{k}}, \hat{X} \tilde{\phi}_{\vec{i}})$.

III. THE EXPECTATION VALUE OF THE METRIC OPERATOR

A. Gaussian weave state

In LQG, the space is constructed by “excitation” of a graph. Actually, if the graph is not include the multivalent vertex that valence is higher than three, the volume of the space is zero. Thus, the “ground state” is not a flat space. In order to obtain a flat space, the graph must include infinite number of vertices. Therefore, the weave state $\mathcal{W} = \prod_{v \in R} w_v$ in finite region R in three-dimensional space is defined as following [30, 33]. Let us consider the two closed edges γ_1, γ_2 crossing at the vertex v . The wave fuction of color- p is

$$\Phi_p = \text{tr}(h_{\gamma_1}^{(p)} h_{\gamma_2}^{(p)}) = (h_{\gamma_1}^{(p)})^A{}_B (h_{\gamma_2}^{(p)})^C{}_D (\phi_0(p, p, p, p))_A{}^B{}_C{}^D. \quad (18)$$

constructed by the edges, as FIG. 2. The inner product of Φ_p is normalized. We take a




FIG. 2: The graph constructed by γ_1, γ_2 . The path of the graph is $\gamma_1^a(0) \rightarrow \gamma_1^a(1) = \gamma_2^a(0) \rightarrow \gamma_2^a(1) = \gamma_1^a(0)$. Four tangent vectors $\dot{\gamma}_1^a(0), \dot{\gamma}_1^a(1), \dot{\gamma}_2^a(0)$ and $\dot{\gamma}_2^a(1)$ are linearly independent. The vertex can take the color-0, 2, $\dots$, $2p$, but we treat the color-0 for simplicity.

weave state for each vertex as $\sum_{p=0}^{\infty} c_p \Phi_p$, and determine the coefficient as following:

$$w_v = N \exp(-\lambda^2(\Phi_1 - 2)^2), \quad (19)$$

where N, λ are normalization factor and any real parameter, respectively. Using the formula of $SU(2)$ tensor products [33]:

$$(\Phi_1)^n = \sum_p \frac{(p+1) \cdot n!}{\left(\frac{n-p}{2}\right)! \left(\frac{n+p}{2} + 1\right)!} \Phi_p, \quad \frac{n-p}{2} = 0, 1, 2, \dots \quad (20)$$

and generating function of Hermite polynomials, $\exp(-t^2 + 2xt) = \sum H_n(x)t^n/n!$, (19) is

$$\begin{aligned} w_v &= N e^{-4\lambda^2} \sum_{n=0}^{\infty} \frac{\lambda^n H_n(2\lambda)}{n!} (\Phi_1)^n \\ &= N e^{-4\lambda^2} \sum_{p=0}^{\infty} \sum_{n=0}^{\infty} \frac{p+1}{n!(n+p+1)!} \lambda^{2n+p} H_{2n+p}(2\lambda) \cdot \Phi_p. \end{aligned} \quad (21)$$

Then we obtain the coefficient as

$$c_p(\lambda) = N e^{-4\lambda^2} \sum_{n=0}^{\infty} \frac{(p+1)\lambda^{2n+p} H_{2n+p}(2\lambda)}{n!(n+p+1)!}. \quad (22)$$

Although the series diverges depending on the value of λ , it is at most $\exp(4\lambda^2)$. However, in numerical point of view, it is preferable to converge it. Moreover, in order to avoid a contribution of higher color, we employ $\lambda = 3/4$ same as [30]. The norm for each vertex as $\langle w_v | w_v \rangle = \sum_{p=0}^{\infty} (c_p(\frac{3}{4}))^2$.

B. Matrix elements of metric operators

(9) is rewritten by the holonomies:

$${}^n\Theta_I^i(v) = -\frac{4}{\kappa n} \text{tr}(\tau^i h_I \{h_I^{-1}, (V_v)^n\}_P), \quad (23)$$

where $h_I = h_{s_I}^{(1)}$, s_I is a segment that endpoint is at v . Let us quantize (23) by usual procedure $\{\cdot, \cdot\}_P \rightarrow \frac{1}{i\hbar}[\cdot, \cdot]$, then

$${}^n\hat{\Theta}_I^i(v) = -\frac{4}{i\hbar\kappa n}\text{tr}(\tau^i h_I[h_I^{-1}, \hat{V}_v^n]) = \frac{4}{i\hbar\kappa n}\text{tr}(\tau^i h_I \hat{V}_v^n h_I^{-1}). \quad (24)$$

${}^n\hat{\Theta}_I^i(v)$ is equal to $\hat{Q}_{e_I}^i(v, n)$ times $-8i\hbar$ defined in [27]. We operate it to w_v . First, h_I^{-1} acts to a trivalent vertex $\tilde{\phi}_k(p, p, p, p)$:

$$\begin{aligned} h_0^{-1} \tilde{\phi}_k(p, p, p, p) &= N_k(p) \cdot 1 \text{ (diagram: a horizontal line with four vertices, each with a vertical line above it labeled } p \text{, and a curved line connecting the first and second vertices from the left)} \\ &= N_k(p) \sum_{q=\pm 1} \alpha_q(p) \cdot 1 \text{ (diagram: a horizontal line with four vertices, each with a vertical line above it labeled } p \text{, and a curved line connecting the first and second vertices from the left, with a label } p+q \text{ below the first vertex and } 1 \text{ below the second vertex)} \end{aligned} \quad (25)$$

where $\alpha_{+1}(p) = 1$, $\alpha_{-1} = -\frac{p}{p+1}$ and $N_k(p) = N_k(p, p, p, p)$. It can be regarded as a pentavalent vertex for volume operator. Denote

$$\tilde{\phi}_{ij}(\check{1}, q, p, p, p) = N_{ij}(\check{1}, q, p, p, p) \text{ (diagram: a horizontal line with five vertices, each with a vertical line above it labeled } p \text{, and a curved line connecting the first and second vertices from the left, with labels } (e_0) q, (e_1) p, (e_2) p, (e_3) p \text{ above the vertices and } i, j \text{ below the first two vertices)} \quad (26)$$

where the check $\check{1}$ means that the volume operator does not operate to it. By (A32), matrix element of volume operator is found, $\hat{V}^n \tilde{\phi}_{ij} = \sum_{s,t} V_{ij}^{n, st} \tilde{\phi}_{st}$, then

$$\begin{aligned} \hat{V}^n h_0^{-1} \tilde{\phi}_k &= \sum_{q=\pm 1} \sum_{s,t} \alpha_q(p) N_k(p) \left(\frac{N_{st} V_{pk}^{n, st}}{N_{pk}} \right) (\check{1}, p+q, p, p, p) \\ &\times 1 \text{ (diagram: a horizontal line with four vertices, each with a vertical line above it labeled } p \text{, and a curved line connecting the first and second vertices from the left, with labels } p+q \text{ below the first vertex and } 1 \text{ below the second vertex)} \end{aligned} \quad (27)$$

Therefore, the action of dreibein operator with density weight $(n-1)$ is obtained as

$$\begin{aligned} {}^n\hat{\Theta}_0^i \tilde{\phi}_k &= -\frac{4}{i\hbar\kappa n} \sum_{q=\pm 1} \sum_{s,t} \alpha_q(p) N_k(p) \left(\frac{N_{st} V_{pk}^{n, st}}{N_{pk}} \right) (\check{1}, p+q, p, p, p) \\ &\times \frac{\text{Tet} \begin{bmatrix} s & p & p+q \\ 1 & 1 & 2 \end{bmatrix}}{\theta(s, p, 2)} \cdot \text{(diagram: a horizontal line with four vertices, each with a vertical line above it labeled } p \text{, and a curved line connecting the first and second vertices from the left, with labels } s, t \text{ below the first two vertices and a loop labeled } i \text{ on the first vertex)} \end{aligned} \quad (28)$$

In right hand side of (28), $su(2)$ generator connects the intertwiner. Since it is not $SU(2)$ invariant, the expectation value becomes zero. Thus we operate it two times, that is, metric operator:

$${}^n\hat{q}(s_I, s_J)(v) = \frac{1}{2}({}^n\hat{\Theta}_I^i(v) {}^n\hat{\Theta}_J^i(v) + {}^n\hat{\Theta}_J^i(v) {}^n\hat{\Theta}_I^i(v)). \quad (29)$$

It operates to two vertices but it is same points, because its expectation value vanishes by spin network calculation when it acts different points. In generally, dreibein operator is non-commutative: $[\hat{\Theta}_I^i(v), \hat{\Theta}_J^j(v)] \neq 0$, thus we take (29) to be symmetric explicitly. Let the graphical part of right hand side of (28) denote as

$$f_{st}^i = \begin{array}{c} \begin{array}{cccc} (e_0) & (e_1) & (e_2) & (e_3) \\ p & p & p & p \end{array} \\ \begin{array}{c} \text{Diagram: A vertex } i \text{ with a loop labeled } 2. \text{ A line labeled } s \text{ goes to a vertex } s. \text{ From } s, \text{ a line labeled } t \text{ goes to a vertex } t. \text{ From } t, \text{ a line labeled } p \text{ goes to a vertex } p. \text{ From } p, \text{ a line labeled } p \text{ goes to a vertex } p. \text{ From } p, \text{ a line labeled } p \text{ goes to a vertex } p. \end{array} \end{array} \quad (30)$$

We operate a dreibein operator to it. There is two cases, $I = J$ and $I \neq J$, in the operation. In the case of $I = J = 0$, the action of h_0^{-1} is

$$h_0^{-1} f_{st}^i = \begin{array}{c} \begin{array}{c} \text{Diagram: A vertex } i \text{ with a loop labeled } 2. \text{ A line labeled } s \text{ goes to a vertex } s. \text{ From } s, \text{ a line labeled } t \text{ goes to a vertex } t. \text{ From } t, \text{ a line labeled } p \text{ goes to a vertex } p. \text{ From } p, \text{ a line labeled } p \text{ goes to a vertex } p. \text{ From } p, \text{ a line labeled } p \text{ goes to a vertex } p. \end{array} \end{array} = \sum_{v=\pm 1} \alpha_v(s) \begin{array}{c} \begin{array}{c} \text{Diagram: A vertex } i \text{ with a loop labeled } 2. \text{ A line labeled } s \text{ goes to a vertex } s. \text{ From } s, \text{ a line labeled } |s+v| \text{ goes to a vertex } |s+v|. \text{ From } |s+v|, \text{ a line labeled } t \text{ goes to a vertex } t. \text{ From } t, \text{ a line labeled } p \text{ goes to a vertex } p. \text{ From } p, \text{ a line labeled } p \text{ goes to a vertex } p. \text{ From } p, \text{ a line labeled } p \text{ goes to a vertex } p. \end{array} \end{array} \quad (31)$$

The volume operator regards it as $\tilde{\phi}_{st}(\check{1}, |s+v|, p, p, p)$. Therefore, the matrix element of the metric operator of same direction is

$$\begin{aligned} {}^n\hat{q}(s_0, s_0)(v) \tilde{\phi}_k &= -\frac{8}{(\hbar\kappa n)^2} \sum_{q=\pm 1} \sum_{s,t} \sum_{v=\pm 1} \sum_m \\ &\times \alpha_q(p) \alpha_v(s) \left(\frac{N_k}{N_m} \right) (p) \\ &\times \left(\frac{N_{st} V_{pk}^{n_{st}}}{N_{pk}} \right) (\check{1}, p+q, p, p, p) \left(\frac{N_{pm} V_{st}^{n_{st}}}{N_{st}} \right) (\check{1}, |s+v|, p, p, p) \\ &\times \frac{\text{Tet} \begin{bmatrix} s & p & p+q \\ 1 & 1 & 2 \end{bmatrix} \text{Tet} \begin{bmatrix} p & s & |s+v| \\ 1 & 1 & 2 \end{bmatrix}}{\Delta_p \theta(s, p, 2)} \tilde{\phi}_m \\ &= \sum_m {}^nq(s_0, s_0)_{(p)km} \tilde{\phi}_m. \end{aligned} \quad (32)$$

Similarly, all ${}^n\hat{q}(s_I, s_J)(v)$ can also be obtained.

Concretely, we show the expectation value of ${}^n\hat{q}(s_I, s_J)$ numerically. In (22), let the norm be normalize $\sum_p (c_p)^2(\lambda) = 1$, and we set a parameter $\lambda = 3/4$, then $(c_0)^2 = 0.414892$, $(c_1)^2 = 0.482013$ and $(c_2)^2 = 0.0972374$. Since the coefficients of higher color more than $p = 3$ are negligible such as $(c_3)^2 = 1.28919 \times 10^{-6}$, it is sufficient to evaluate the contribution of $p = 1, 2$. Then we obtain

$$\begin{aligned} {}^1q(s_0, s_0)_{00} &= 0.150\hbar\kappa, \\ {}^1q(s_1, s_1)_{00} &= 0.131\hbar\kappa, \\ {}^1q(s_2, s_2)_{00} &= 0.265\hbar\kappa, \\ {}^1q(s_3, s_3)_{00} &= 0.0980\hbar\kappa. \end{aligned} \tag{33}$$

The state (18) is not symmetric with respect to edges, the values of length squared of tangent vectors at v are different each other. We can assign a distance between vertices to an average of $\dot{s}_I$'s length, $\frac{1}{4} \sum_I \sqrt{{}^1q(s_I, s_I)_{00}} = 0.394(\hbar\kappa)^{1/2}$. The some angles between $\dot{s}_I^a$'s, which defined by $\theta(s_I, s_J) = \cos^{-1}({}^1q(s_I, s_J)_{00} / (\sqrt{{}^1q(s_I, s_I)_{00}} \sqrt{{}^1q(s_J, s_J)_{00}})$, are also obtained as

$$\begin{aligned} \theta(s_0, s_1) &= 160^\circ, \\ \theta(s_0, s_2) &= 74.6^\circ, \\ \theta(s_0, s_3) &= 83.0^\circ. \end{aligned} \tag{34}$$

Thus, the direction of $\dot{s}_I^a$ is not symmetric in this state.

IV. CONCLUSION

We calculated that the expectation value of the metric operator with respect to Gaussian weave state. It can be identified to the length squared of the edge $|\dot{s}_I^a|^2$, and it determines the scale of this system. We found that the diagonal element of the metric is surely positive definite. This result seems non-trivial at first sight, because the metric operator

$${}^n\hat{q}(s_I, s_I) = \frac{4}{(\hbar\kappa n)^2} \left(2\text{tr}(h_I \hat{V}^{2n} h_I^{-1}) - \left(\text{tr}(h_I \hat{V}^n h_I^{-1}) \right)^2 \right) \tag{35}$$

has a negative sign in second term, and the matrix element of the volume operator is not positive definite. (The eigenvalue of it is positive definite.) However, by the spin network

calculation,

$$\begin{aligned}
||^n \hat{\Theta}_0^i \tilde{\phi}_k||^2 &= \sum_{s,t} (\text{positive number}) \times \text{Diagram} \\
&= -(\text{positive number}) \times \text{tr}(\tau^i \tau^i) \geq 0, \quad (\text{no-sum with respect to } i). \quad (36)
\end{aligned}$$

The diagram is a tensor network with two external indices \$i\$ and \$i\$ on the left, each with a loop labeled 2. These are connected to a central box with four vertical lines. The top two lines are labeled \$s\$ and \$t\$, and the bottom two are labeled \$s\$ and \$t\$. The box is divided into three vertical sections, each labeled \$p\$.

That is, $\langle w_v | ^n \hat{q}(s_I, s_I) w_v \rangle = \sum_i ||^n \hat{\Theta}_I^i w_v||^2 \geq 0$.

Acknowledgments

I would like to thank Ken-Ichi Aoki, for helpful comments and useful advice.

APPENDIX A: SPIN NETWORK CALCULATION

1. Spin network and recoupling theory

In this subsection, we referred to [5]. The $SU(2)$ invariant tensor, a matrix X^A_B and its trace are graphically written as

$$\begin{aligned}
\delta_B^A &= \begin{array}{c} A \\ | \\ B \end{array}, \quad \delta_D^A \delta_C^B = - \begin{array}{c} A \quad B \\ \diagdown \quad \diagup \\ C \quad D \end{array}, \quad i\epsilon^{AB} = \begin{array}{c} A \quad B \\ \cup \end{array} \\
i\epsilon_{AB} &= \begin{array}{c} A \quad B \\ \cap \end{array}, \quad X^A_B = \begin{array}{c} A \\ \boxed{X} \\ B \end{array}, \quad \text{tr} X = - \begin{array}{c} \boxed{X} \end{array}. \quad (A1)
\end{aligned}$$

Since we add the minus signature to the crossing line, a symmetric tensor becomes anti-symmetric line. We derive the binor identity

$$\begin{array}{c} | \\ | \end{array} + \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} + \begin{array}{c} \cup \\ \cap \end{array} = 0. \quad (A2)$$

Thus, the lines behave as knots satisfying the Reidemeister moves O, I, II and III:

$$\begin{aligned}
\text{O} : & \quad \begin{array}{c} \cap \\ \cup \end{array} = \begin{array}{c} | \\ | \end{array}, \\
\text{I} : & \quad \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} = \begin{array}{c} \cap \\ \cap \end{array}, \\
\text{II} : & \quad \begin{array}{c} \cap \\ \cap \end{array} = \begin{array}{c} \cup \\ \cup \end{array}, \\
\text{III} : & \quad \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} = \begin{array}{c} \diagdown \quad \diagup \\ \diagup \quad \diagdown \end{array} \quad (A3)
\end{aligned}$$

in the knot theory if they are fixed the indices upper or lower position. A line of color- a is a pieces of lines that anti-symmetrized:

$$\left| \begin{smallmatrix} \cdots \\ \cdots \end{smallmatrix} \right|_a = \underbrace{\left| \begin{smallmatrix} \cdots \\ \cdots \end{smallmatrix} \right|}_a = \frac{1}{a!} \left(\left| \begin{smallmatrix} \cdots \\ \cdots \end{smallmatrix} \right| - \left| \begin{smallmatrix} \cdots \\ \cdots \end{smallmatrix} \right| + \cdots \right), \quad (\text{A4})$$

where the white box means anti-symmetrizing of lines. A trivalent vertex defined as

$$\begin{array}{c} a \\ | \\ b \text{---} \bullet \text{---} c \end{array} = \begin{array}{c} a \\ | \\ \text{---} \text{---} \text{---} \\ | \quad | \quad | \\ b \quad i \quad c \end{array}, \quad \begin{cases} i = \frac{1}{2}(-a + b + c) \\ j = \frac{1}{2}(a - b + c) \\ k = \frac{1}{2}(a + b - c) \end{cases}. \quad (\text{A5})$$

The generator of $su(2)$ anti-hermite 2-dimensional representation is

$$\tau^i = \begin{array}{c} | \\ \circlearrowleft i \end{array} = \begin{array}{c} \circlearrowleft i \end{array} \begin{array}{c} 2 \\ | \end{array}. \quad (\text{A6})$$

This generator acts to a line of color- a as

$$\begin{array}{c} \circlearrowleft i \end{array} \begin{array}{c} 2 \\ | \end{array} \left| \begin{smallmatrix} \cdots \\ \cdots \end{smallmatrix} \right|_a = a \cdot \begin{array}{c} \circlearrowleft i \end{array} \begin{array}{c} 2 \\ | \end{array} \left| \begin{smallmatrix} \cdots \\ \cdots \end{smallmatrix} \right|_a. \quad (\text{A7})$$

Summation of product of generators

$$\sum_{i=1}^3 \begin{array}{c} \circlearrowleft i \end{array} \begin{array}{c} 2 \\ | \end{array} = \frac{1}{2} \cdot \begin{array}{c} 2 \\ \text{---} \end{array}, \quad \sum_{i,j,k} \epsilon^{ijk} \begin{array}{c} \circlearrowleft i \end{array} \begin{array}{c} 2 \\ | \end{array} \begin{array}{c} \circlearrowleft j \end{array} \begin{array}{c} 2 \\ | \end{array} \begin{array}{c} \circlearrowleft k \end{array} \begin{array}{c} 2 \\ | \end{array} = -\frac{1}{2} \cdot \begin{array}{c} 2 \\ \text{---} \end{array}. \quad (\text{A8})$$

The symmetrizer

$$\Delta_a = \bigcup^a = (-1)^a (a+1). \quad (\text{A9})$$

The θ -net

$$\theta(a, b, c) = \begin{array}{c} a \\ \bullet \text{---} \bullet \\ b \\ \bullet \text{---} \bullet \\ c \end{array} = \frac{(-1)^{m+n+p} (m+n+p+1)! m! n! p!}{a! b! c!}, \quad (\text{A10})$$

where $m = \frac{1}{2}(-a + b + c)$, $n = \frac{1}{2}(a - b + c)$ and $p = \frac{1}{2}(a + b - c)$. The tetrahedral net

$$\text{Tet} \begin{bmatrix} a & b & e \\ c & d & f \end{bmatrix} = \begin{array}{c} b \quad c \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \\ a \quad d \\ f \end{array} \begin{array}{c} e \\ | \end{array} = \frac{\mathcal{I}}{\mathcal{E}} \sum_{m \leq S \leq M} \frac{(-1)^S (S+1)!}{\prod_i (S - a_i)! \cdot \prod_i (b_i - S)!}, \quad (\text{A11})$$

$$\begin{aligned}
a_1 &= (a + d + e)/2, \quad b_1 = (b + d + e + f)/2, \\
a_2 &= (b + c + e)/2, \quad b_2 = (a + c + e + f)/2, \\
a_3 &= (a + b + f)/2, \quad b_3 = (a + b + c + d)/2, \\
a_4 &= (c + d + f)/2, \\
m &= \max\{a_i\} \quad , \quad M = \min\{b_i\} \\
\mathcal{E} &= a!b!c!d!e!f! \quad , \quad \mathcal{I} = \prod_{ij}(b_j - a_i)!.
\end{aligned}$$

The 9- j symbol

$$\left\{ \begin{array}{ccc} a & b & c \\ d & e & f \\ g & h & i \end{array} \right\} = \text{Diagram} \quad . \quad (\text{A12})$$

The exchanging of lines in a trivalent vertex

$$\text{Diagram} = \lambda_c^{ab} \text{Diagram} \quad , \quad \lambda_c^{ab} = (-1)^{\frac{1}{2}(a(a+3)+b(b+3)-c(c+3))}. \quad (\text{A13})$$

The recoupling theorem

$$\text{Diagram} = \sum_e \left\{ \begin{array}{ccc} a & b & e \\ c & d & f \end{array} \right\} \text{Diagram} \quad . \quad (\text{A14})$$

The relation between a 6- j symbol and a tetrahedral net

$$\left\{ \begin{array}{ccc} a & b & e \\ c & d & f \end{array} \right\} = \frac{\Delta_e}{\theta(a, d, e)\theta(b, c, e)} \text{Tet} \left[\begin{array}{ccc} a & b & e \\ c & d & f \end{array} \right]. \quad (\text{A15})$$

The reduction formulas

$$\text{Diagram} = \delta_{ad} \frac{\theta(a, b, c)}{\Delta_a} \Big|_a, \quad (\text{A16})$$

$$\text{Diagram} = \frac{\text{Tet} \left[\begin{array}{ccc} a & b & e \\ c & d & f \end{array} \right]}{\theta(a, f, b)} \text{Diagram} \quad . \quad (\text{A17})$$

2. Matrix elements of volume operator

The volume operator for the vertex v is

$$\hat{V} = \frac{(\hbar\kappa)^{3/2}}{4} \sqrt{\sum_{I < J < K} |i\hat{W}_{[IJK]}|} \quad (\text{A18})$$

where $|\hat{X}| = \sqrt{\hat{X}^\dagger \hat{X}}$. The meaning of the square root is that its diagonalized matrix equivalent to square root of $\hat{X}^\dagger \hat{X}$. The summation I, J, K is the label of edges e_I connected with the vertex v . The operator $\hat{W}_{[IJK]}$ adding the three $su(2)$ generators:

$$\hat{W}_{[IJK]} = \frac{1}{2} \left(\overset{(e_I)}{\underset{2}{\curvearrowright}} + \overset{(e_J)}{\underset{2}{\curvearrowright}} + \overset{(e_K)}{\underset{2}{\curvearrowright}} \right) \quad (\text{A19})$$

(times minus two) to the edges e_I, e_J, e_K (see (A8)). Let us compute this operation to the n -valent vertex. Its matrix element is

$$\begin{aligned} \widetilde{W}_{[IJK]} \vec{i}^{\vec{k}} &= (\tilde{\phi}_{\vec{k}}, \hat{W}_{[IJK]} \tilde{\phi}_{\vec{i}}) \\ &= N_{\vec{i}} N_{\vec{k}} P_I P_J P_K \cdot P_0 \left(\text{Diagram} \right) P_{n-1}. \end{aligned} \quad (\text{A20})$$

The diagram in (A20) is a graph with three vertical strands. The left strand has vertices labeled $i_I, i_{I+1}, \dots, i_{I+1}$ and i_{I+1} at the bottom. The middle strand has vertices labeled $i_J, i_{J+1}, \dots, i_{J+1}$ and i_{J+1} at the bottom. The right strand has vertices labeled $i_K, i_{K+1}, \dots, i_{K+1}$ and i_{K+1} at the bottom. The strands are connected by horizontal lines labeled P_I, P_J, P_K and P_I, P_J, P_K respectively. The bottom of the diagram is labeled $2, 2, 2$.

We can calculate the graphical part of (A20). Now we separate it four part (i) e_0-e_I , (ii) e_I-e_J , (iii) e_J-e_K and (iv) e_K-e_{n-1} .

(i)

$$\begin{aligned} P_0 \left(\text{Diagram} \right) &= \lambda_{P_I}^{P_I^2} \cdot P_0 \left(\text{Diagram} \right) \\ &= -\lambda_{k_{I+1}}^{i_{I+1}^2} \left(\prod_{x=2}^r \delta_{i_x}^{k_x} \right) \frac{\theta(P_0, P_1, i_2)}{\Delta_{i_2}} \left(\prod_{x=2}^{r-1} \frac{\theta(i_x, P_x, i_{x+1})}{\Delta_{i_{x+1}}} \right) \\ &\quad \times \frac{\text{Tet} \begin{bmatrix} k_{I+1} & i_{I+1} & i_I \\ P_I & P_I & 2 \end{bmatrix}}{\theta(k_{I+1}, i_{I+1}, 2)} \cdot \left(\text{Diagram} \right) \end{aligned} \quad (\text{A21})$$

The diagram in (A21) is a graph with two vertical strands. The left strand has vertices labeled $i_I, i_{I+1}, \dots, i_{I+1}$ and i_{I+1} at the bottom. The right strand has vertices labeled i_{I+1} and i_{I+1} at the bottom. The strands are connected by horizontal lines labeled P_I and P_I respectively. The bottom of the diagram is labeled $2, 2$.

$$= \alpha_i \cdot \begin{array}{c} \text{---} k_{I+1} \\ \text{---} \bullet \\ \text{---} i_{I+1} \\ \text{---} 2 \end{array} \quad (\text{A22})$$

(iv)

$$\begin{aligned} \left(\begin{array}{c} k_K \text{---} k_{K+1} \\ \text{---} P_K \\ 2 \text{---} P_K \\ \text{---} i_K \text{---} i_{K+1} \end{array} \right) P_{n-1} &= \left(\prod_{x=t+1}^{n-2} \delta_{i_x}^{k_x} \right) \left(\prod_{x=t+1}^{n-3} \frac{\theta(i_x, P_x, i_{x+1})}{\Delta_{i_x}} \right) \frac{\theta(i_{n-2}, P_{n-2}, P_{n-1})}{\Delta_{i_{n-2}}} \\ &\times \frac{\text{Tet} \begin{bmatrix} i_K & k_K & i_{K+1} \\ P_K & P_K & 2 \end{bmatrix}}{\theta(k_K, i_K, 2)} \cdot \begin{array}{c} k_K \text{---} \\ 2 \text{---} \bullet \\ i_K \text{---} \end{array} \end{aligned} \quad (\text{A23})$$

$$= \alpha_{iv} \cdot \begin{array}{c} k_K \text{---} \\ 2 \text{---} \bullet \\ i_K \text{---} \end{array} \quad (\text{A24})$$

Then the graph of (A20) is

$$P_0 \left(\begin{array}{c} k_I \text{---} k_{I+1} \quad k_J \text{---} k_{J+1} \quad k_K \text{---} k_{K+1} \\ \text{---} P_I \quad \text{---} P_J \quad \text{---} P_K \\ \text{---} P_I \quad \text{---} P_J \quad \text{---} P_K \\ i_I \text{---} i_{I+1} \quad i_J \text{---} i_{J+1} \quad i_K \text{---} i_{K+1} \\ \text{---} 2 \quad \text{---} 2 \quad \text{---} 2 \end{array} \right) P_{n-1} = \alpha_i \alpha_{iv} \cdot \begin{array}{c} k_{I+1} \text{---} k_J \text{---} k_{J+1} \text{---} k_K \\ \text{---} \bullet \quad \text{---} \bullet \quad \text{---} \bullet \\ \text{---} i_{I+1} \text{---} i_J \text{---} i_{J+1} \text{---} i_K \\ \text{---} 2 \quad \text{---} 2 \quad \text{---} 2 \end{array} \quad (\text{A25})$$

Next, we calculate the remaining parts (ii) and (iii).

(ii)

$$\begin{aligned} \begin{array}{c} k_{I+1} \text{---} k_{J-1} \text{---} k_J \\ 2 \text{---} \bullet \quad \text{---} P_{s-1} \\ i_{I+1} \text{---} i_{J-1} \text{---} i_J \end{array} &= \left(\prod_{x=r+1}^{s-1} \frac{\text{Tet} \begin{bmatrix} k_x & k_{x+1} & 2 \\ i_{x+1} & i_x & P_x \end{bmatrix}}{\theta(k_{x+1}, i_{x+1}, 2)} \right) \begin{array}{c} k_J \\ 2 \text{---} \bullet \\ i_J \end{array} \end{aligned} \quad (\text{A26})$$

$$= \alpha_{ii} \cdot \begin{array}{c} k_J \\ 2 \text{---} \bullet \\ i_J \end{array} \quad (\text{A27})$$

(iii)

$$\begin{aligned}
& \begin{array}{c} k_{J+1} \quad k_{J+2} \\ \text{---} \text{---} \\ | \quad | \\ P_{s+1} \\ | \quad | \\ i_{J+1} \quad i_{J+2} \\ \text{---} \text{---} \\ 2 \end{array} \begin{array}{c} \curvearrowright \\ k_K \\ \curvearrowleft \end{array} i_K = \lambda_{k_K}^{i_K 2} \cdot \begin{array}{c} k_{J+1} \quad k_{J+2} \\ \text{---} \text{---} \\ | \quad | \\ P_{s+1} \\ | \quad | \\ i_{J+1} \quad i_{J+2} \\ \text{---} \text{---} \\ 2 \end{array} \begin{array}{c} \curvearrowright \\ k_K \\ \curvearrowleft \end{array} i_K \\
&= \frac{\lambda_{k_K}^{i_K 2}}{\lambda_{k_{J+1}}^{i_{J+1} 2}} \left(\prod_{x=s+1}^{t-1} \frac{\text{Tet} \begin{bmatrix} k_x & k_{x+1} & 2 \\ i_{x+1} & i_x & P_x \end{bmatrix}}{\theta(k_x, i_x, 2)} \right) \begin{array}{c} k_{J+1} \\ \text{---} \\ | \\ i_{J+1} \\ \text{---} \\ 2 \end{array} \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array}
\end{aligned} \tag{A28}$$

$$= \alpha_{\text{iii}} \cdot \begin{array}{c} k_{J+1} \\ \text{---} \\ | \\ i_{J+1} \\ \text{---} \\ 2 \end{array} \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} \tag{A29}$$

Then we obtain

$$\begin{aligned}
\widetilde{W}_{[IJK]i}^{\vec{k}} &= N_i N_{\vec{k}} P_I P_J P_K \cdot P_0 \left(\begin{array}{c} k_I \quad k_{I+1} \quad k_J \quad k_{J+1} \quad k_K \quad k_{K+1} \\ \text{---} \text{---} \text{---} \text{---} \text{---} \text{---} \\ | \quad | \quad | \quad | \quad | \quad | \\ P_I \quad P_J \quad P_K \\ | \quad | \quad | \quad | \quad | \quad | \\ i_I \quad i_{I+1} \quad i_J \quad i_{J+1} \quad i_K \quad i_{K+1} \\ \text{---} \text{---} \text{---} \text{---} \text{---} \text{---} \\ 2 \quad 2 \end{array} \right) P_{n-1} \\
&= N_i N_{\vec{k}} P_I P_J P_K \cdot \alpha_i \alpha_{\text{ii}} \alpha_{\text{iii}} \alpha_{\text{iv}} \cdot \left\{ \begin{array}{ccc} k_J & P_J & k_{J+1} \\ i_J & P_J & i_{J+1} \\ 2 & 2 & 2 \end{array} \right\}.
\end{aligned} \tag{A30}$$

Since $i\widetilde{W}_{[IJK]}$ is a pure imaginally anti-symmetric matrix, it can be diagonalized. Using unitary matrix $U_{[IJK]}$ which diagonalize it,

$$\sum_{I < J < K} |i\hat{W}_{[IJK]}| \tilde{\phi}_{\vec{i}} = \sum_{I < J < K} \sum_{\vec{j}, \vec{k}} U_{[IJK]}^\dagger \vec{i} \tilde{j} |i\widetilde{W}_{[IJK]}^{\text{diag}}| U_{[IJK]} \vec{j} \tilde{k} \tilde{\phi}_{\vec{k}} \tag{A31}$$

The sum of hermite matrices is also hermite, (A18) can be diagonalized. Thus, the matrix element of $\hat{V}^n$ is

$$\begin{aligned}
\hat{V}^n \tilde{\phi}_{\vec{i}} &= \sum_{\vec{j}, \vec{k}} R_i^{\dagger \vec{j}} ((V^2)_{\vec{j}}^{\text{diag}})^{n/2} R_{\vec{j}}^{\vec{k}} \tilde{\phi}_{\vec{k}} \\
&= \frac{(\hbar\kappa)^{3n/2}}{4^n} \sum_{\vec{j}, \vec{k}, \vec{l}, \vec{m}, \vec{n}} R_i^{\dagger \vec{j}} [R_{\vec{j}}^{\vec{k}}] \sum_{I < J < K} U_{[IJK]}^\dagger \vec{l} \tilde{i} |i\widetilde{W}_{[IJK]}^{\text{diag}}| U_{[IJK]} \vec{l} \tilde{m} [R_{\vec{m}}^{\dagger \vec{j}}]^{n/2} R_{\vec{j}}^{\vec{n}} \tilde{\phi}_{\vec{n}}.
\end{aligned} \tag{A32}$$

-
- [1] C. Rovelli, Living Rev. Rel. **1**, 1 (1998), gr-qc/9710008.
 - [2] C. Rovelli and L. Smolin, Nucl. Phys. **B442**, 593 (1995), gr-qc/9411005.
 - [3] R. Loll, Phys. Rev. Lett. **75**, 3048 (1995), gr-qc/9506014.
 - [4] S. Frittelli, L. Lehner, and C. Rovelli, Class. Quant. Grav. **13**, 2921 (1996), gr-qc/9608043.
 - [5] R. De Pietri and C. Rovelli, Phys. Rev. **D54**, 2664 (1996), gr-qc/9602023.
 - [6] A. Ashtekar and J. Lewandowski, Adv. Theor. Math. Phys. **1**, 388 (1998), gr-qc/9711031.
 - [7] C. Rovelli, Phys. Rev. Lett. **77**, 3288 (1996), gr-qc/9603063.
 - [8] A. Ashtekar, A. Corichi, and J. A. Zapata, Class. Quant. Grav. **15**, 2955 (1998), gr-qc/9806041.
 - [9] R. Borissov, R. De Pietri, and C. Rovelli, Class. Quant. Grav. **14**, 2793 (1997), gr-qc/9703090.
 - [10] M. Gaul and C. Rovelli, Class. Quant. Grav. **18**, 1593 (2001), gr-qc/0011106.
 - [11] M. Takeda et al., Phys. Rev. Lett. **81**, 1163 (1998), astro-ph/9807193.
 - [12] N. Hayashida et al., Astrophys. J. **522**, 225 (1999), astro-ph/0008102.
 - [13] K. Greisen, Phys. Rev. Lett. **16**, 748 (1966).
 - [14] G. T. Zatsepin and V. A. Kuzmin, JETP Lett. **4**, 78 (1966).
 - [15] G. Amelino-Camelia, J. R. Ellis, N. E. Mavromatos, D. V. Nanopoulos, and S. Sarkar, Nature **393**, 763 (1998), astro-ph/9712103.
 - [16] J. R. Ellis, N. E. Mavromatos, and D. V. Nanopoulos, Gen. Rel. Grav. **31**, 1257 (1999), gr-qc/9905048.
 - [17] H. Sato and T. Tati, Prog. Theor. Phys. **47**, 1788 (1972).
 - [18] R. Gambini and J. Pullin, Phys. Rev. **D59**, 124021 (1999), gr-qc/9809038.
 - [19] J. Alfaro, H. A. Morales-Tecotl, and L. F. Urrutia, Phys. Rev. **D65**, 103509 (2002), hep-th/0108061.
 - [20] L. Urrutia, Mod. Phys. Lett. **A17**, 943 (2002), gr-qc/0205103.
 - [21] J. Alfaro, H. A. Morales-Tecotl, and L. F. Urrutia, Phys. Rev. Lett. **84**, 2318 (2000), gr-qc/9909079.
 - [22] P. Jain, D. W. McKay, S. Panda, and J. P. Ralston, Phys. Lett. **B484**, 267 (2000), hep-ph/0001031.
 - [23] H. Davoudiasl, J. L. Hewett, and T. G. Rizzo, Phys. Lett. **B549**, 267 (2002), hep-ph/0010066.
 - [24] K. Hagiwara and Y. Uehara, Phys. Lett. **B517**, 383 (2001), hep-ph/0106320.

- [25] A. Ibarra and R. Toldra, JHEP **06**, 006 (2002), hep-ph/0202111.
- [26] L. A. Anchordoqui, J. L. Feng, and H. Goldberg, Phys. Lett. **B535**, 302 (2002), hep-ph/0202124.
- [27] H. Sahlmann and T. Thiemann (2002), gr-qc/0207030.
- [28] H. Sahlmann and T. Thiemann (2002), gr-qc/0207031.
- [29] T. Thiemann, Class. Quant. Grav. **18**, 2025 (2001), hep-th/0005233.
- [30] A. Corichi and J. M. Reyes, Int. J. Mod. Phys. **D10**, 325 (2001), gr-qc/0006067.
- [31] T. Thiemann, Class. Quant. Grav. **15**, 839 (1998), gr-qc/9606089.
- [32] T. Thiemann, Class. Quant. Grav. **15**, 1281 (1998), gr-qc/9705019.
- [33] M. Arnsdorf and S. Gupta, Nucl. Phys. **B577**, 529 (2000), gr-qc/9909053.